

## ORIGINAL ARTICLE

## EFFECTS OF NATURAL HONEY ON BLOOD GLUCOSE AND LIPID PROFILE IN YOUNG HEALTHY PAKISTANI MALES

Muhammad Majid, M. Azeem Younis, Abdul Khaliq Naveed\*, Muhammad Usman Shah, Zahid Azeem\*, Syed Haider Tirmizi

Final year MBBS Student, \*Department of Biochemistry and Molecular Biology, Army Medical College, National University of Sciences and Technology, Rawalpindi, Pakistan

**Background:** Honey has been shown to have beneficial effects on glucose and lipid profiles in patients at high risk of heart diseases. Therefore, this study was carried out to investigate the effects of natural honey on blood glucose and lipid profile in healthy individuals. **Methods:** A randomized controlled trial was carried out in the Army Medical College, Rawalpindi, Pakistan, spanning 4 weeks, that is, from 15<sup>th</sup> February to 15<sup>th</sup> March 2009. A total of 70 healthy young boarders of the same college were included in the study and randomly divided into two groups of 35 each using random number table. Seventy gram (70g) of honey was given to each individual in the experimental group daily for a period of 4 weeks while control group was kept on the same diet as that of experimental group except honey. The fasting glucose, total cholesterol, low density lipoprotein (LDL), high density lipoprotein (HDL) and triglyceride (TG) levels were measured before and after the experiment. **Results:** Fasting glucose levels in both groups were raised. However, the increase in the experimental group was significantly less than that in the control group ( $p < 0.05$ ). The levels of total cholesterol, LDL and triglycerides in the control group increased while those in the experiment group decreased significantly ( $p < 0.05$ ), while HDL levels were decreased in the former and increased in the latter group ( $p < 0.05$ ). **Conclusion:** Natural honey consumption significantly limits the rise in blood glucose along with a significant decrease in the levels of total cholesterol, LDL and triglycerides, and increase HDL in young healthy adults.

**Keywords:** Honey, Glucose profile, Lipid profile.

J Ayub Med Coll Abbottabad 2014;26(1):42-5

## INTRODUCTION

From ancient times, honey is known to be one of the most beneficial drugs of nature. Through the ages people have found many uses of it. It was used as an alternative to gold by the Romans, in matrimonial ceremonies by the Greeks, and in the treatment of wounds by the Egyptians.<sup>1,2</sup> Honey has got dynamic and multidimensional medicinal and surgical uses. In a research it is established that honey improves insulin sensitivity.<sup>3</sup> Significant increase in the insulin secretion capacity is associated with decrease in circulating leptin, total cholesterol, and LDL.<sup>4</sup> Cardiac risk factors, such as obesity, smoking, hypertension, and chronic periodontal disease, are associated with elevated C-reactive protein (CRP) levels and this marker was decreased by antioxidants present in honey.<sup>5,6</sup> In an experiment on dogs, it has been noted that fructose is having a unique ability to increase uptake of hepatic glucose and insulin that is suggestive of its modulating property of glycaemic control.<sup>7</sup> Fructose being a major component of honey improves the glycaemic response to a glucose load.<sup>8,9</sup> The harmful and genotoxic effects of mycotoxins are reduced and the gut microflora is improved by honey.<sup>10</sup> Honey is helpful in the treatment of psoriasis<sup>11</sup>, diaper dermatitis<sup>12</sup>, *Pityriasis versicolor*, *Tinea cruris*, *Tinea corporis* and *Tinea faciei*.<sup>13</sup> Honey is very effective in healing of wounds.<sup>14</sup> Honey

increases the production against thymus-dependent and thymus-independent antigens in primary and secondary immune responses.<sup>15</sup> Natural honey lowers plasma prostaglandin concentrations.<sup>16</sup> Honey improves haematological indices.<sup>17</sup> Recent researches by Al-Waili and Chepulis have shown that natural honey lowers fasting blood glucose, total cholesterol, LDL, VLDL, TG's and increases HDL, thus reducing cardiovascular risks.<sup>5,6</sup>

Since honey shows these effects by multiple proposed mechanisms which might be influenced by the genotypes, geographical distribution of the subjects and the type of honey used, so the effects of honey collected from Pakistan were seen on Pakistani subjects in our study. Age, gender, diet (except honey) and life style of the subjects which may confound the results were kept constant in our study. The objective of this study was to investigate the effects of natural honey collected from Pakistan on blood glucose and lipid profile in healthy Pakistani individuals keeping strict control conditions including same sex, place of living, life style, diet (except honey) and more or less same age of the subjects.

## MATERIAL AND METHODS

This randomized controlled trial (RCT) was carried out in Army Medical College, Rawalpindi, Pakistan from 15<sup>th</sup> Feb to 15<sup>th</sup> March 2009. Before the start of study,

Research Advisory Committee approved the research and informed written consent was taken from all subjects. The subjects of this research were newly admitted first year students of Army Medical College, Rawalpindi, Pakistan and their complete physical and clinical examination and laboratory investigations were done which is mandatory for admission to the college. Their results showed that all had normal complete blood and cardiac function. The subjects were in the same hostel on the same diets and the same daily routine. Seventy subjects were included in the study through convenient sampling and randomly divided into two groups of 35 each using random number table. Seventy grams (70g) of honey was given daily for a period of four weeks to each individual in the experimental group dissolved in 250 ml of tap water and then ingested while control group was kept on the same diet as that of experimental group except honey. Honey was purchased from Ilyas Traders, Charsadda, Khyber Pakhtunkhwa, Pakistan. The honeybees (*Apis mellifera*) fed on trees of *Acacia modesta*. The honey was natural and unprocessed. Three subjects from experimental group and four subjects from control group dropped out because their daily routine and diet changed significantly during the course of study and some of them started taking medicines which could affect the results of our study. The fasting glucose, total cholesterol, LDL, HDL, TG levels were studied at the start of the study and then after 4 weeks. Laboratory investigations were performed using spectra 2 auto-analyzer, by Merck Company, Germany.

Data was analysed using SPSS-15. Quantitative variables were described as mean±standard deviation (SD). Paired sample's *t*-test was applied to compare initial and final levels of study variables within each group. Change in study variables were compared between the groups through independent sample's *t*-test. A *p*-value of ≤0.05 was taken as significant.

## RESULTS

Mean age of the boys of experimental group was 20.13±0.14 years and of control group was 20±0.15 years. Both the groups were comparable with respect to age (*p*=0.537), fasting blood glucose (*p*=0.083), total cholesterol (*p*=0.060), TG (*p*=0.195), LDL (*p*=0.137) and HDL level (*p*=0.142).

In experimental group, significant decrease was observed in total cholesterol (*p*<0.001) and LDL level (*p*=0.001) while HDL level was increased significantly (*p*=0.005) while change in glucose and triglyceride were not significant. In control group, significant increase was observed in glucose (*p*<0.001), total cholesterol (*p*<0.001), triglyceride (*p*=0.009) and LDL (*p*=0.002) while insignificant decrease was observed in HDL level (*p*=0.433) as tabulated in Table-1.

The changes in all the variables within four weeks between both the groups were compared. Increase in serum fasting glucose level in experimental group was significantly lower than that of control group (*p*=0.011). Total cholesterol was decreased in experimental group while it was increased in the control group and this change was significantly different between both the groups (*p*<0.001). The TG level was reduced in experimental group where as it increased in control group and this difference in change was significant (*p*=0.018). In LDL level decrease in experimental group and an increase in control group was observed and the change between both the groups was significantly different (*p*<0.001). The HDL level was increased in experimental and decreased in control group. Change in mHDL level in experimental group was significantly different as compared to that in control group (*p*=0.013). (Table-2)

**Table-1: Within the group comparison of initial and final levels**

Variables (mmol/L)	Control (n=31)		Cases (n=32)	
	Initial	Final	Initial	Final
Glucose	4.88±0.06	5.27±0.06*	5.03±0.06	5.16±0.08 <sup>NS</sup>
Total cholesterol	3.54±0.10	3.83±0.13*	3.89±0.15	3.49±0.11*
TG	0.70±0.04	0.84±0.04*	0.82±0.09	0.76±0.05 <sup>NS</sup>
LDL	2.58±0.08	2.81±0.11*	2.78±0.10	2.49±0.08*
HDL	0.722±0.02	0.706±0.02 <sup>NS</sup>	0.779±0.03	0.823±0.03*

Data was expressed as mean±SE, \* Final level is significantly different from initial level. <sup>NS</sup> Final level is insignificantly different from initial level

**Table-2: Comparison of change between both the groups**

Variables (mmol/L)	Control (n=31)	Cases (n=32)	<i>p</i> -value
Glucose	0.390±0.074	0.122±0.071	0.011*
Total cholesterol	0.287±0.052	0.403±0.092	<0.001*
TG	0.139±0.050	0.062±0.060	0.018*
LDL	0.228±0.066	0.288±0.075	<0.001*
HDL	-0.016±0.020	0.044±0.014	0.013*

Data was expressed as mean±SE, \*Levels are significantly different between both the groups

## DISCUSSION

In the present study we measured serum levels of glucose, triglycerides, total cholesterol, LDL and HDL. The levels of these substances are regulated within an optimum range by body homeostatic mechanisms. But their levels are subject to change by various internal and external environmental factors. One of the significant factors is the dietary changes that the modern lifestyle has brought along with it in the form of fast foods. Changes in the levels of these substances make the human body prone to various clinical disorders. Glucose is a monosaccharide, the immediate source of energy for human body and its level is increased in Diabetes Mellitus. In the past cholesterol had been portrayed by some a poison but it has dichotomy of good (HDL) and

bad (LDL) and is essential for the synthesis of bile acid and some of steroid hormones.<sup>18</sup> Triglycerides are major component of VLDL and plays key role in metabolism as energy source and in fat transport but its role in cardiac risk factor is also reported.<sup>5,6</sup> LDL is also involved in transportation of cholesterol to different body tissues where only small amount is used and rest of other is free.<sup>19</sup> Thus, LDL plays a part in the development of cardiac complications. Elevated level of native LDL is not associated with cardiac risk rather oxidized LDL is involved in the formation of atherosclerotic plaque.<sup>20</sup> HDL is a good cholesterol, reservoir of apolipo-proteins, take up the unesterified cholesterol both from other lipoprotein particles and from cell membranes and then esterify it and takes it to liver, thus helping in reverse cholesterol transport.<sup>19</sup> Elevated levels of HDL decreases cardiovascular risk factors.<sup>5</sup> In our study, the effects of honey on the plasma levels of these various bio-molecules have been recorded. Glucose levels in both groups were raised. However, increase in the fasting glucose levels of the experimental group was significantly less than those in the control group. Levels of total cholesterol, LDL and triglycerides in the control group increased while those in the experiment group decreased significantly. HDL level was increased in experimental group whereas it decreased slightly in control group and this difference was significant. This substantiates the work done by other researchers.<sup>5,6,21</sup> Honey alters the plasma levels of these substances by various biochemical mechanisms. Honey has been shown to decrease blood glucose level. Honey has got stimulatory effect on the secretion of insulin and also improves insulin sensitivity<sup>3</sup>, thereby reducing glucose level. Honey produces hydrogen peroxide which has insulin like effects.<sup>21</sup> Honey has nitric oxide (NO) metabolites and probably stimulates NO synthase and so increases NO production.<sup>22</sup> Nitric oxide is stimulatory to the release of insulin.<sup>23</sup> It has been found that honey decreases the plasma and urinary levels of some prostaglandins like PGE<sub>2</sub>, PGF<sub>2</sub> and TXB<sub>2</sub>.<sup>16,22</sup> Prostaglandins are inhibitory to insulin secretion.<sup>24</sup> Zinc and Copper are normal constituents of honey. So administration of honey increases serum zinc and copper levels which play significant role in the metabolism of insulin and glucose.<sup>17,25</sup> Fructose in the honey may decrease the hyperglycaemic response of glucose content of honey.<sup>5</sup> It has been found that low dose fructose administered with glucose decreased the glycaemic response to a glucose load in healthy individuals and type 2 diabetic patients.<sup>8,9</sup> It was proposed that fructose acted possibly by stimulating glucokinase translocation.<sup>9</sup> So honey might stimulate glucokinase to take up the glucose into the liver.<sup>5</sup> Honey decreased TGs, total cholesterol and LDL. As discussed earlier honey stimulates insulin release. Insulin is stimulatory to lipoprotein lipase which cause breakdown

of triglycerides present in plasma lipoproteins to free fatty acids and glycerol, which are transported to peculiar sites and get metabolized.<sup>19</sup> Administration of large amounts of fructose increases TGs, total cholesterol and LDL.<sup>26-28</sup> Administration of low dose fructose along with glucose causes the opposite effect; they decreased TG, total cholesterol and LDL.<sup>27,28</sup> The same effect has been achieved with honey. It has also been found that fructose in smaller concentrations decrease hepatic cholesterol synthesis by decreasing the activity of hepatic 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase.<sup>29</sup> Niacin is a normal constituent of honey although present in lesser amount. Lipolysis in adipose tissue results primarily in increase level of circulating free fatty acids in blood which are taken up by liver and used in TGs synthesis.<sup>19</sup> Niacin strongly inhibits this lipolysis in adipose tissue which results in decrease in hepatic TGs synthesis and thus TGs plasma levels.<sup>19</sup> TGs synthesis is required for VLDL synthesis and LDL is derived from VLDL in blood plasma.<sup>19</sup> So niacin decreases plasma TGs, total cholesterol and LDL. Honey might mediate a part of its lowering effect on TGs, total cholesterol and LDL through niacin. As mentioned earlier oxidized LDL is responsible for atherosclerotic plaque formation. Honey decreases oxidized LDL due to presence of anti-oxidants in honey like ascorbic acid, pyridoxine, pantothenic acid, riboflavin and Thiamine. In the present research increase in HDL in experimental group is significant as compared to the decrease in control group. As discussed earlier, HDL gets cholesterol from cell membranes and other lipoproteins like LDL and transports it to the liver. We have also seen that honey decreases LDL levels. So, less HDL is used up in transporting cholesterol from LDL to liver because of decreased LDL. Like this less LDL is used up in its physiologic process, so HDL levels rises. So honey increase HDL levels indirectly by decreasing LDL, this needs further experimentation. Foods with high glycaemic index have been associated with decreased HDL levels.<sup>30</sup> So Honey being a low glycaemic index food increases HDL.<sup>5</sup> Increase in HDL levels has been associated with weight loss and nicotinic acid.<sup>31</sup> Honey reduces weight and so increases HDL levels.<sup>5</sup> Nicotinic acid (niacin) present in honey although in lesser amounts might increases HDL levels.

## CONCLUSION

Consumption of honey for a period of 4 weeks is effective in reducing glucose, TG's, total cholesterol, LDL and increasing HDL in young healthy adults. Therefore, healthy individuals should include honey in their diet to improve their glucose and lipid profile, and to prevent acquiring diseases in which the levels of glucose, TGs, total cholesterol and LDL is increased and

HDL is decreased like diabetes, cardiovascular diseases, hyperlipidaemias, and obesity.

## ACKNOWLEDGEMENTS

We are thankful to National University of Science and Technology (NUST) for funding our research project. We are also thankful to Maj. Dr. M. Jawad Yousaf from Department of Biochemistry and Molecular Biology, AMC, Rawalpindi and Rafay Ali Sabzwari from AMC, Rawalpindi who helped us a lot in our project. We are indebted to Miss Irum Abid who is biostatistician in PAFMJ office, AMC, Rawalpindi for her effort in carrying out statistics of this project. We present great appreciation to the staff of Pathology laboratory, Military Hospital Rawalpindi for assisting us in lab procedures of our research.

A grant of Rs. 30000 was received by the authors from the National University of Sciences and Technology, Pakistan, as part of its undergraduate research promotion program.

## REFERENCES

- Crane E, editor. The archaeology of beekeeping. London: Duckworth; 1983.
- Pliny the Elder, editor. Naturalis Historica. Rome: Hackios; 1669.
- Katsilambros NL, Philippides P, Toulitau A, Georgakopoulos K, Kofotzouli L, Frangaki D, *et al.* Metabolic effects of honey (alone or combined with other foods) in type II diabetics. *Acta Diabetologica Latina* 1988;25:197–203.
- Cuhadaroglu C, Utkusavas A, Ozturk L, Salman S, Ece T. Effects of Nasal CPAP treatment on insulin resistance, lipid profile, and plasma leptin in sleep apnea. *Lung* 2009;187(2):75–81.
- Yaghoobi N, Al-Waili N, Ghayour-Mobarhan M, Parizadeh SM, Abasali Z, Yaghoobi Z, *et al.* Natural honey and cardiovascular risk factors; Effects on Blood Glucose, Cholesterol, Triacylglycerole, CRP and Body weight Compared with Sucrose. *Sci World J* 2008;8:463–9.
- Al-Waili, NS. Natural honey lowers plasma glucose, C-reactive protein, homocysteine, and blood lipids in healthy, diabetic, and hyperlipidemic subjects: comparison with dextrose and sucrose. *J Med Food* 2004;1:100–7.
- Watford M. Small amounts of dietary fructose dramatically increase hepatic glucose uptake through a novel mechanism of glucokinase activation. *Nutr Rev* 2002;60:253–7.
- Moore MC, Davis SN, Mann SL, Cherrington AD. Acute fructose administration improves oral glucose tolerance in adults with type 2 diabetes. *Diabetes Care* 2001;24:1882–7.
- Moore MC, Cherrington AD, Mann SL, Davis SN. Acute fructose administration decreases glycaemic response to an oral glucose tolerance test in normal adults. *J Clin Endocrinol Metab* 2000;85:4515–9.
- Ezz El-Arab AM, Girgis SM, Hegazy EM, Abd El-Khalek AB. Effect of dietary honey on intestinal microflora and toxicity of mycotoxins in mice. *BMC Complement Altern Med* 2006;6:6.
- Al-Waili, NS. Topical application of natural honey, beeswax and olive oil mixture for atopic dermatitis or psoriasis: partially controlled, single-blinded study. *Complement Ther Med* 2003;11:226–34.
- Al-Waili NS. Clinical and mycological benefits of topical application of honey, olive oil and beeswax in diaper dermatitis. *Clin Microbiol Infect* 2005;11:160–3.
- Al-Waili NS. An alternative treatment for pityriasis versicolor, tinea cruris, tinea corporis and tinea faciei with topical application of honey, olive oil and beeswax mixture: an open pilot study. *Complement Ther Med* 2004;12:45–7.
- Okany CC, Atimomo CE, Akinyanju OO. Efficacy of natural honey in the healing of leg ulcers in sickle cell anaemia. *Niger Postgrad Med J* 2004;11:179–81.
- Al-Waili NS, Haq A. Effect of honey on antibody production against thymus-dependent and thymus-independent antigens in primary and secondary immune responses. *J Med Food* 2004;7:491–4.
- Al-Waili, N NS, Boni NS. Natural honey lowers plasma prostaglandin concentrations in normal individuals. *J Med Food* 2003;6:129–33.
- Al-Waili NS. Effects of daily consumption of honey solution on hematological indices and blood levels of minerals and enzymes in normal individuals. *J Med Food* 2003;6:135–40.
- Hume R, Boyd GS. Cholesterol metabolism and steroid hormone production. *Biochem Soc Trans* 1978;6:893–8.
- Champe PC, Harvey RA, Ferrier DR, editors. Lippincott's Illustrated Reviews: Biochemistry. 4<sup>th</sup> ed. New Delhi: Wolters Kluwer; 2008.
- MacDonald-Wicks L, Garg M. Oxidized LDL and Antioxidants in Atherosclerosis. In: Cheema SK, editor. *Biochemistry of atherosclerosis*. Australia: Springer; 2006;p.519.
- Chepulis L, Starkey N. The long-Term Effects of Feeding Honey Compared with Sucrose and a Sugar-Free Diet on Weight Gain, Lipid Profiles, and DEXA Measurements in Rats. *J Food Sci* 2007;73(1):H1–7.
- Al-Waili NS. Effects of honey on the urinary total nitrite and prostaglandins concentration. *Int Urol Nephrol* 2005;37:107–11.
- Smukler SR, Tang L, Wheeler MB, Salapatek AM. Exogenous nitric oxide and endogenous glucose-stimulated beta cell nitric oxide augment insulin release. *Diabetes* 2002;51:3450–60.
- Tran PO, Gleason CE, Piotout V, Robertson RP. Prostaglandin E2 mediates inhibition of insulin secretion by interleukin-1 beta. *J Biol Chem* 1999;274:31245–8.
- Marreiro DN, Geloneze B, Tambascia MA, Lerário AC, Halpern A, Cozzolino SM. Effect of zinc supplementation on serum leptin levels and insulin resistance of obese women. *Biol Trace Elem Res* 2006;112(2):109–18.
- Bantle JP, Swanson JE, Thomas W, Laine DC. Metabolic effects of dietary fructose in diabetic subjects. *Diabetes Care* 1992;15:1468–76.
- Abraha A, Humphreys SM, Clark ML, Mathews DR, Frayn, KN. Acute effect of fructose on postprandial lipaemia in diabetic and non-diabetic subjects. *Br J Nutr* 1998;80:169–75.
- Swanson JE, Laine DC, Thomas W, Bantle JP. Metabolic effects of dietary fructose in healthy subjects. *Am J Clin Nutr* 1992;55:851–6.
- Feingold KR, Moser AH. Effect of glucose or fructose feeding on cholesterol synthesis in diabetic Animals. *Am J Physiol Gastrointest Liver Physiol* 1985;249(5 pt 1):G634–41.
- Frost G, Leeds A, Dore C, Madeiros S, Branding S, Dornhorst A. Glycemic index as a determinant of serum HDL-cholesterol concentration. *Lancet* 1999;353:1045–8.
- Drexel H. Reducing risk by raising HDL-cholesterol: the evidence. *Eur Heart J* 2006;8 (Suppl F):F23–9.

## Address for Correspondence:

Muhammad Usman Shah, House No. 516, Street 5, Sector C4, Phase 5, Hayatabad, Peshawar, Pakistan.

Cell: +92-346-9187704

Email: usman\_shah44@msn.com