

# ADOLESCENT SELF REPORTED REPRODUCTIVE MORBIDITY AND HEALTH CARE SEEKING BEHAVIOUR

M. Mizanur Rahman, M. Kabir, M. Shahidullah\*

Department of Statistics, Jahangirnagar University, Savar, Dhaka and \*Department of Population Dynamics, National Institute of Preventive and Social Medicine, Mohakhali, Dhaka

**Background:** Addressing reproductive health issues of women is now on the global social agenda in the new millennium. Maternal mortality has long been the only indicator of women's health even though reproductive morbidity occurs far more frequently and seriously affects women's lives. In this paper, an attempt was made to assess the magnitude of self reported gynaecological morbidity unrelated to childbearing among the adolescents irrespective of their marital status. The paper also explored the determinants of health care seeking behaviour of the adolescents for their reproductive ailments. **Methods:** Both quantitative and qualitative data were collected for this study. Cross sectional study was conducted both in rural and urban areas using a multistage cluster sampling technique. A nationally representative data on 2883 adolescents irrespective of their marital status were analysed. **Results:** Analysis revealed that a large proportion of the adolescents (64.5%) reportedly has been suffering from gynaecological morbidity. The most frequent form of morbidity was menstrual disorders (63.9%) followed by lower abdominal pain (58.6%), burning sensation during urination(46.1%), genital itching (15.5%), vaginal discharge (3.4%) etc. Multivariate logistic regression analysis revealed that older adolescents aged 15-19 years, family income, type of family, type of residence and hygienic practice during menstruation appeared to be influencing factors for adolescents reproductive morbidity. The results also revealed that about one fifth (18.0%) sought health care for their gynaecological ailments indicating that adolescents were unaware about their reproductive morbidity ( $p < 0.05$ ). For assessing the factors influencing their health care seeking for reproductive morbidity, multivariate logistic regression analysis found significant positive association with adolescents aged 15-19 years, having autonomy in treatment, working status, adolescents of joint or extended family ( $p < 0.05$ ). **Discussion:** Adolescent reproductive health in Bangladesh indicates high incidence of maternal morbidity. The incidence of these health problems varied by socio-economic and demographic characteristics of the adolescents. Recommendations to address these problems include encouragement of female education, introduction of family life education in school curricula, creating community awareness for seeking health care and empowerment of women in household decision making process.

**Key Words:** Adolescents, reproductive morbidity, care seeking behaviour, Bangladesh.

## INTRODUCTION

In developing countries, reproductive morbidity commonly affects the quality of women's lives. This form of ill health has largely been ignored by the policy makers, health planners as well as researchers. The reproductive morbidity includes the obstetric and gynaecological conditions of ill health related to the reproductive process during and outside the childbearing episodes. The obstetric morbidity encompasses the conditions during pregnancy, delivery and post partum period and gynaecological morbidity includes the conditions outside pregnancy related events<sup>1</sup>. In this part, the reproductive morbidity refers to gynaecological morbidity of ill health unrelated to pregnancy.

There are three methods for the diagnosis of gynaecological morbidity such as self reported symptoms, clinical examination and laboratory tests. Appropriate laboratory testing is considered as the "Gold standard" for the precise detection of reproductive morbidity and accurate measurement of prevalence of diseases. However, such tests have limited applicability in developing countries because they are expensive and present logistical difficulties<sup>2</sup>. Health facilities at the community level are poorly equipped to deal with reproductive morbidity, since they do not have diagnostic facilities, drugs, supply of blood or surgical equipment to treat the diseases. Even, service providers are not well acquainted to detect the morbidity or to provide counselling.

Information about reproductive morbidity in developing countries is scanty. Although a few studies have been conducted in this field, but most of them are based on information obtained from clinics or hospitals. Large proportion of women does not visit health facilities unless the disease becomes serious. So, the results from hospitals or clinics do not reflect the magnitude of the disease burden. The statistics provided by the hospitals are based on biomedical causes only, but information on social, economic, demographic and behavioural determinants are rare. In fact, a search of literature reveals that knowledge about reproductive morbidity and its determinants in Bangladesh and also in the sub-continent are almost non-existent. A few studies in this area showed a varying prevalence of reproductive morbidity<sup>3-6</sup> and these mainly considered the adult women of reproductive age. Adolescent reproductive health has been ignored. Concerted efforts are needed to provide useful information for health planners and policy makers. So, the appropriate strategies to be designed to bring about improvement in the reproductive health of women<sup>5</sup>. With these objectives in view, the present study on self reported reproductive morbidity among adolescents was undertaken.

## MATERIALS AND METHODS

This was a cross sectional study conducted in both rural and urban areas of Bangladesh. The female adolescents aged 10-19 years irrespective of their marital status constituted the study population. A multistage cluster sampling technique was adopted to select the sample. Both quantitative and qualitative data on reproductive morbidity were collected. A total of 2883 adolescents were selected. They were also inquired about their health care seeking behaviour for their problems. Both uni-variate and bi-variate analyses were performed. Multivariate logistic regression analysis was used to identify the socio-economic and demographic factors which were significantly related to reproductive morbidity and care seeking behaviour. Data analysis was performed using Statistical Package for Social Science (SPSS, version 11.0). To substantiate the results of quantitative study, a series of focus group discussions were also conducted among the adolescents. In the present study, the analysis on quantitative data was presented.

## RESULTS

### Socio-demographic characteristics

The mean age of the respondents was  $16.3 \pm 1.8$  years with a range of 10 to 19 years. Among the respondents, 34.9% were unmarried and 65.1% were married. The mean years of schooling was  $4.7 \pm 3.3$  years. Among them 23.6% were illiterate, 33.1% had 1-5 years of schooling and the rest had 6 and above years of schooling. Regarding parental education, more than two fifths (43.7%) of the adolescents' fathers were illiterate as against more than two thirds (69.9%) of mothers were illiterate indicating that mothers were more illiterate than fathers. The mean family size was 5.2 persons. About one third (29.6%) of the adolescents were currently engaged in different income generating activities other than household work. Overwhelmingly majority of the respondents were Muslims (89.0%) and only 11.0% were non-Muslims. About three fifths of the adolescents (57.3%) were from nuclear families and the rest from joint/ or extended families (42.7%). The median family income was Tk.2500.0 and about three fifths (58.1%) of the families had income less than Tk. 3000.0 (Table 1).


**Table –1: Socio-demographic characteristics of the adolescents (N=2883)**

			<b>Mean</b>
<b>Characteristics</b>	<b>Frequency</b>	<b>%</b>	<b>± SD</b>
<b>Age in years</b>			
10-14	567	19.7	16.3±1.8
15-19	2316	80.3	
<b>Marital status</b>			
Unmarried	1005	34.9	
Married	1878	65.1	
<b>Residence</b>			
Rural	1743	60.5	

Urban	1140	39.5	
<b>Years of schooling (Res)</b>			
0	680	23.6	4.7±3.3
1-5	955	33.1	
≥6	1248	43.3	
<b>Religion</b>			
Non-Muslim	317	11.0	
Muslim	2566	89.0	
<b>Level of education (F)</b>			
Illiterate	1260	43.7	
Literate	1623	56.3	
<b>Level of education (M)</b>			
Illiterate	2016	69.9	
Literate	867	30.1	
<b>Work status</b>			
No	2030	70.4	
Yes	853	29.6	
<b>Type of family</b>			
Nuclear	1651	57.3	
Joint	1232	42.7	
<b>Family size</b>			
2-3	862	23.7	5.2±2.1
4-5	1050	36.4	
≥6	1151	39.9	
<b>Monthly family income (Tk.)</b>			
<2000	569	23.0	Median income=
2000-3000	871	35.1	Tk. 2500.0
3000-4000	609	24.6	
≥4000	430	17.3	

### Overall reproductive morbidity and frequency

Married adolescents with current pregnancy and unmarried adolescents not menstruating (that is who did not attain the age of menarche) were excluded from the analysis. So, a total of 2883 adolescents were studied. The adolescents were inquired about symptoms of gynaecological morbidity for the last six months. Among them 35.5% had no gynaecological problem and the rest 64.5% had one or more gynaecological morbidities. The average number of problems was 1.9 ranging from 1-6. Two-fifths (40.0%) had one problem followed by 36.4% who had two, 18.6% have had three and 5.0% had four and more problems. The most frequent problem was menstrual disorders (63.9%) followed by lower abdominal pain (58.6%), burning urination (46.1%), genital itching (15.5%), vaginal discharge with fever (3.4%), genital ulcer (1.6%). More than one-fourth of the married adolescents (26.4%) had complain of pain during sexual intercourse and 1.9% had complain of bleeding after sexual intercourse (Married adolescents only) (Fig-1).



**Figure-1: Frequency of self reporting morbidity**

#### **Variation of self reported reproductive morbidity by marital status and age**

Analysis revealed that the reproductive morbidities were found significantly higher among married than unmarried adolescents such as genital itching was 2.8 times, vaginal discharge 2.4 times, burning urination 1.6 times and lower abdominal pain 1.5 times higher among the married adolescents whereas the menstrual disorders were less among the married adolescents ( $p < 0.05$ ). On the other hand, the reproductive morbidities were significantly high among the older adolescents aged 15-19 years except pain during sexual intercourse among the married adolescents ( $p < 0.05$ ). Genital itching was 3.1 times, vaginal discharges 2.4 times, lower abdominal pain 1.9 times, burning urination 1.8 times and menstrual disorders 1.4 times higher among the older adolescents. But older married adolescents were less likely to complain of painful sexual intercourse. (This might be due to immaturity of physical development of younger adolescents). On the other hand, the reproductive morbidities were higher among older and married adolescents. This might be due to their longer exposure to sexual environment.

#### **Correlates of self reported symptoms of morbidity: Multivariate analysis**

The present study found that 64.5% of the adolescents reported about one or more gynaecological problems. This indicates a high rate of reproductive morbidity among the adolescents. To examine the socio-demographic and behavioural factors associated with reproductive morbidity, the variables which were found to be significantly associated in bi-variate analysis were considered in the logistic regression analysis. The dependent variable was dichotomous i.e. whether the adolescents reported any gynaecological problem in the last six months. Adolescents' current age was found to be significantly associated with reported morbidities. Older adolescents reported 1.8 times more gynaecological morbidity than younger adolescents. Similarly, adolescents having monthly family income Tk. 4000.0 and above 1.6 times and having monthly income Tk. 2000.0-3000.0 1.3 times reported about the symptoms of morbidity. The adolescents of joint or extended families were 1.4 times likely to report symptoms of morbidity. But adolescents of urban background and hygienic practice during menstruation were less likely to report about gynaecological morbidity.

#### **Care Seeking Behaviour for reported morbidity**

### Type of health facilities attended by adolescents:

Adolescents attended institutional as well as non-institutional health care facilities for their gynaecological morbidities. About sixty five per cent of the adolescents who had complain of gynaecological problems, only 18.0% attended health care facilities for treatment.

Although majority of the adolescents attended institutional health care facilities, a large proportion of them preferred to take treatment from quacks (25.4%). More than one tenth of the adolescents (11.9%) received treatment from traditional healers such as Homeopaths or Kabirajes (practitioners of indigenous medicine). About institutional health care receivers, most of them received treatment from Thana Health Complexes ( 26.5%) followed by Family Welfare Centres (16.7%) and from other government hospitals1(2.8%).

**Reasons for not receiving treatment:** The major reasons, as mentioned by adolescents for not receiving any health care for gynaecological problems were personal grounds which include, 41.0% mentioned ‘no need of treatment’ followed by ‘lack of knowledge’ 19.8%, ‘economic hardship’ 18.4% and ‘shyness to expose to doctor’ 14.8%. In addition 6.1% of the adolescents reasoned related to inadequate service facilities such as ‘no female doctor available in the hospital’.

### Determinants of care seeking behaviour for reproductive morbidity: Multi-variate analysis

In order to examine the contribution of each of the factors and health care seeking behaviour, a multivariate logistic regression analysis was performed with dichotomous dependent variable “whether received health care or not”. Analysis revealed that older adolescents aged 15-19 years were 3.4 times more likely to seek health care than their younger counterparts. Most importantly, adolescents having autonomy for their treatment that is having control over other members of the family were 1.4 times more likely to seek health care for their reproductive problems. Analysis also revealed that working adolescents and members of joint or extended families were 1.3 times more likely to seek health care.

## DISCUSSION

Most of the previous studies ignored the reproductive morbidities of unmarried adolescents, but unmarried adolescents are also exposed to similar environment as of married adolescents with the exception of marital sex. Older women were included in the study of Bhatia et al.,<sup>5</sup> and Wasscrheit et al<sup>3</sup>.

**Table 2: Type of self reported reproductive morbidity by marital status**

Reproductive morbidity	N=2883	Marital status		Odds ratio	95% CI
		Married n=1878			
<b>Unmarried n=1005</b>					
<i>Menstrual disorders</i>					
Yes	41.2	38.3	46.5	0.716***	0.613-0.836
No	58.8	61.7	53.5		
<i>Lower abdominal pain</i>					
Yes	37.8	40.9	31.8	1.484***	1.263-1.744
No					

62.2	59.1	68.2			
<b><i>Burning urination</i></b>					
Yes	29.7	33.0	23.7	1.584***	1.331-1886
No	70.3	67.0	76.3		
<b><i>Genital itching</i></b>					
Yes	10.0	12.7	5.0	2.785***	2.032-3.817
No	90.0	87.3	95.0		
<b><i>Vaginal discharge</i></b>					
Yes	2.2	2.8	1.2	2.357***	1.252-4.435
No	97.8	97.2	98.8		
<b><i>Genital ulcer</i></b>					
Yes	1.0	1.2	0.7	1.768	0.756-4.134
No	99.0	98.8	99.3		
<b><i>*Pain during sexual intercourse</i></b>					
Yes	17.2	26.4	-	-	
No	82.8	73.6	100.0		

<b>*Bleeding after sexual intercourse</b>					
Yes	1.2	1.9	0.0	-	-
No	98.8	98.1	100.0		

\*\*\* $p < 0.001$ , Reference category = Unmarried adolescents, \*Married adolescents

**Table 3: Type of self reported reproductive morbidity by age**

Reproductive morbidity	Total N=2883	Age in years		Odds ratio	95% CI
		10-14 (n=567)			
<b>15-19 (n=2316)</b>					
<b>Menstrual disorders</b>					
Yes	41.2	34.9	42.7	1.3889***	1.471-1.6817
No	58.8	65.1	57.3		
<b>Burning urination</b>					
Yes	29.7	21.0	31.9	1.7607***	1.412-2.1944
No	70.3	79.0	68.1		
<b>Lower abdominal pain</b>					
Yes	37.8	26.1	40.6	1.9375***	1.5785-2.3782
No	62.2	73.9	59.4		
<b>Genital itching</b>					
Yes	10.0	4.1	11.5	3.0690***	1.9838-4.7478
No	90.0	95.9	88.5		
<b>Vaginal discharge</b>					
Yes	2.2	1.1	2.5	2.4013**	1.0309-5.5931
No	97.8	98.9	97.5		
<b>Genital ulcer</b>					
Yes	1.0	0.4	1.2	3.4560	0.8211-14.5470
No	99.0	99.6	98.8		
<b>*Pain during sexual intercourse (n=1878)</b>					
Yes	26.4	24.5	53.3	0.2842***	0.1954-0.4134
No	73.6	46.7	75.5		
<b>*Bleeding after sexual intercourse (n=1878)</b>					
Yes	1.9	3.3	1.8	0.5377	0.1870-1.561
No	98.1	98.2	96.7		

\*\*\* $p < 0.001$ ; \*\* $p < 0.01$  Reference category = 10-14 years, \*Married adolescents

**Table 4: Adolescent self reported reproductive morbidity: Multi-variate analysis**

Attributes	$\beta$	<i>p</i> value	Odds ratio	95% CI
<b>Age in years</b>				
10-14 (RC)	-	-	-	-
15-19	0.6062	0.0000	1.8335	1.4341-2.3441
<b>Marital status</b>				
Unmarried (RC)	-	-	-	-
Married	-0.1390	0.2316	0.8702	0.6930-1.0928
<b>Residence</b>				
Rural (RC)	-	-	-	-
Urban	-0.2882	0.0024	0.7496	0.6224-.9029
<b>Level of education (M)</b>				
Illiterate (RC)	-	-	-	-
Literate	-0.1813	0.0739	0.8342	0.6838-1.0176
<b>Monthly family income (Tk.)</b>				
<2000 (RC)	-	-	-	-
2000-3000	0.2400	0.0380	1.2712	1.0134-1.5948
3000-4000	0.1519	0.2274	1.1641	0.9096-1.4897
≥4000	0.5002	0.0007	1.6490	1.2351-2.2017
<b>Work status</b>				
No (RC)	-	-	-	-
Yes	0.1553	0.1068	1.1681	0.9671-1.4108
<b>Type of family</b>				
Nuclear (RC)	-	-	-	-
Joint/extended	0.3676	0.0001	1.4443	1.2077-1.7273
<b>Practice of personal hygiene</b>				
Non sanitary (RC)	-	-	-	-
Sanitary	-0.8530	0.0000	0.4261	0.3020-.6014
Model chi square	104.672			
Df	10			
Significance	0.0000			
N	2479			
Constant	0.638			

\*RC= Reference category

\*\*Variables not included in the regression model are: religion, years of schooling, father's level of education and family size, i.e. these are not statistically significant in bi-variate analysis

**Table 5: Adolescent care seeking behaviour for reproductive morbidity: Multi-variate analysis**

Attributes	$\beta$	<i>p</i> value	Odds ratio	95% CI
<b>Age in years</b>				
10-14 (RC)	-	-	-	-
15-19	1.2331	0.0000	3.4318	2.0694-5.6910
<b>Marital status</b>				
Unmarried (RC)	-	-	-	-
Married	-0.0943	0.5300	0.9100	.6780-1.2214
<b>Working status</b>				
No (RC)	-	-	-	-
Yes	0.2904	0.0246	1.3370	1.0379-1.7224
<b>Type of family</b>				
Nuclear (RC)	-	-	-	-
Joint/extended	0.2782	0.0259	1.3208	1.0340-1.6871
<b>Autonomy for treatment</b>				
No (RC)	-	-	-	-
Yes	0.3581	0.0170	1.4307	1.0661-1.9199
Model chi square	57.008			
df	5			
Significance	0.0000			
N	1859			
Constant	-2.8569			



\*RC= Reference category, \*\*Variables not included in the regression model are: residence, religion, years of schooling, father's and mother's level of education, monthly family income and family size, i.e. these are not statistically significant in bi-variate analysis

Therefore in this study, unmarried adolescents were also included to find out the variation of their reproductive morbidity. In recent years, attention has focused on the problem of high level of maternal mortality in developing countries. Little is known about the prevalence of reproductive morbidity. Dixon-Mueller et al.,<sup>7</sup> opined of "culture of silence" surrounding women's health that typifies these countries and the constraints of living conditions particularly for poor women that prevent the use of health services.

Varying proportion of reproductive morbidity was reported by different studies ranging from 22.0% to 92.0%<sup>3,4,8</sup>. The study revealed a heavy disease burden of reproductive morbidity. The vast majority of the adolescents have been suffering from at least one reproductive morbidity. This high prevalence of reproductive morbidity is certainly disabling women in the community under study, who are mostly illiterate having low socio-economic status. This high prevalence of morbidity raises great concern about women's physical and social well being which causes physical discomfort, personal embarrassment, marital discord and also problems of women's ability to achieve a sustained marital satisfaction<sup>3</sup>. Bang et al<sup>4</sup>., in rural India surprisingly found a high prevalence of gynaecological morbidity of 92 per cent. Both men and women in developing countries with low level of education and lack of health information, especially about personal hygiene could promote misconceptions about many illnesses and limit preventive practices that lead to increased reproductive morbidity. Women of these areas tend to internalize their health problems because of their status in the family, they may not be allowed to seek health care, or they may feel shy about reporting such sort of reproductive problems causing them to be stigmatized by the community<sup>9,10</sup>. Another important factor might hinder them from seeking health care due to inadequate facilities under the government health infrastructure and or inaccessibility of public hospitals. Most of the public hospitals run with inadequate logistics and manpower in the relevant areas of maternal health care.

Analysis of the study revealed an important programme aspect of reproductive health. The older adolescents with rural background, joint or extended family and non-hygienic practice during menstruation are the causes of high prevalence of reproductive morbidity. Similarly, older adolescents having earnings and autonomy in treatment encouraged them to seek health care from any source. This is consistent with South Indian women<sup>5</sup>. So, the programme of health education should be designed in the line of women status in the family.

In conclusion, the study recommends for creating community awareness about health care facilities and instil self concern in adolescents for their own health needs. The first referral units at the grass root levels are Thana Health Complexes and Family Welfare Centres are required to be equipped infrastructurally as well as with skilled manpower for addressing reproductive health problems of women and to provide appropriate referral services. Built in service component and confidentiality may improve self reporting of reproductive morbidity in rapid health surveys. Studies have shown that self reporting is close to clinical diagnosis when diagnostic criterion is clear<sup>11</sup>. Thus such surveys could be an inexpensive way for generating continuous information on reproductive health issues for health managers. The improved and clear diagnostic algorithm for reproductive morbidity help in treating the women with reproductive ailments that will be more fruitful and cost effective in the context of socio-cultural milieu of Bangladesh.

## **ACKNOWLEDGEMENT**

The research was funded by a grant from Social Science Research Council (SSRC), Ministry of Planning, Government of the Peoples Republic of Bangladesh. We acknowledge Director, SSRC for financial assistance.

## **REFERENCES**

1. Zurayk H, Khatib H, Younis N, El-Mouelhy M, Fadle M. Concepts and measures of reproductive morbidity. *Health Transition Review* 1993;3(1):17-40.

2. Jain A, Stein K, Arends-Kuenning M, Garate MR. Measuring reproductive morbidity through a sample survey in Peru. In: IUSSP Seminar in Manila, Philippines, on Innovative Approaches to the Assessment of Reproductive Health. IUSSP and Population Institute, University of the Philippines, Liege, Belgium. 1996.
3. Wassrheit Judith N, Jeffery RH, Chakraborty J, Bradford AK. Reproductive Tract Infections in a family planning population in rural Bangladesh. *Studies in Family Planning* 1989;20(2):69-80.
4. Bang RA, Bang AT, Batulc M, Choudhury Y. High Prevalence of Gynaecological diseases in rural Indian women. *Lancet* 1989;1:85-8.
5. Bhatia JC, Cleland J. Self reported symptoms of gynaecological morbidity and their treatment in South India. *Studies in Family Planning* 1995; 26(4):203-16.
6. Bangladesh Demographic and Health Survey, 1999-2000. National Institute of Population Research and Training (NIPORT), Dhaka, Bangladesh. 2001.
7. Dixon-Mueller, Ruth, Wasserheit J. *The Culture of Silence: Reproductive Tract Infections among women in the third world*. New York: International Health Women's Health Coalition. 1991.
8. Kambo IP, Dhillon BS, Singh P, Saxena BN, Saxena NC. Self reported gynaecological problems from twenty three districts of India. *Indian Journal of Community Medicine* 2003; 27(2):67-73.
9. Khattab H. *The Silent Endurance: Social conditions of women's reproductive Health in Rural Egypt*. Amman: UNICEF and Cairo: The Population Council. 1992.
10. Barua A. 2000. Reproductive Health needs of Married Adolescent Girls in Rural Maharashtra. Paper presented at National Workshop on Reproductive Health. <http://www.rhgateway.com/archive/21.pdf>. Date: 10/14/2003.
11. Zurayk H, Khattab H, Younis N, El-Mouelhy M, Fadle M. Comparing women's reports with medical diagnosis of reproductive morbidity conditions in rural Egypt. *Studies in Family Planning* 1995;26(1):14-21.

---

**Address for Correspondence:**

**Dr. M. Mizanur Rahman**, 58/K, West Raza Bazar, Indira Road, Tejgaon, Dhaka-1215, BANGLADESH.

**Email:** [aniq@citechco.net](mailto:aniq@citechco.net)